


03-29-00  
Express Mail Label No.: EL 430 716 648 US

A  
PATENT



## NYDEGGER & ASSOCIATES

ATTORNEYS AT LAW

348 OLIVE STREET  
SAN DIEGO, CALIFORNIA 92103

NEIL K. NYDEGGER

email@nydegger.com

TELEPHONE  
(619) 688-1300  
FACSIMILE  
(619) 688-1322

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Box Patent Application  
Assistant Commissioner for Patents  
Washington, D.C. 20231

Sir:

Transmitted herewith for filing is the Continuation-in-Part patent application of Tibor Juhasz and J. Randy Alexander for A METHOD OF CORNEAL SURGERY BY LASER INCISING A CONTOURED CORNEAL FLAP comprising sixteen (16) pages of specification and claims.

Also enclosed are:

- X   four ( 4 ) sheets of drawings.
- X   A Certificate of Mailing by "Express Mail" certifying a filing date of March 27<sup>th</sup>, 2000 by use of Express Mail Label No. EL 430 716 648 US.
- X   No fees are enclosed.
- X   No Declaration, Power of Attorney and Petition is Enclosed.
- X   Any additional extension of time required for the timely submission of this paper, the fees for which have not been previously paid, is hereby petitioned and requested.

Please address all future correspondence in connection with the above-identified patent application to the attention of the undersigned.

Dated this 27<sup>th</sup> day of March, 2000.

Respectfully submitted,

Neil K. Hyleger

NEIL K. NYDEGGER  
Attorney for Applicant  
Registration No. 30,202

**NYDEGGER & ASSOCIATES**  
348 Olive Street  
San Diego, California 92103  
Telephone: (619) 688-1300

applwithmsptspto:ddb  
Docket: 11236.11

PATENT

CERTIFICATE OF MAILING BY "EXPRESS MAIL"

"Express Mail" Mailing Label No.: EL 430 716 648 US

Date of Deposit: March 27, 2000

I hereby certify that this utility Patent Application, including claims and drawings, in the names of Tibor Juhasz and J. Randy Alexander, for A METHOD OF CORNEAL SURGERY BY LASER INCISING A CONTOURED CORNEAL FLAP, is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. § 1.10 on the date indicated above in an envelope addressed to Box Patent Application, Assistant Commissioner for Patents, Washington, D.C. 20231.

Dated this 27<sup>th</sup> day of March, 2000.

Respectfully submitted,



DEBRA D. BURNS  
Legal Assistant

NYDEGGER & ASSOCIATES  
348 Olive Street  
San Diego, California 92103  
Telephone: (619) 688-1300

expmailcert  
Docket: 11236.11

[illegible][illegible]

	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2
--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	---

[illegible]

This Application is a continuation-in-part of Application Serial No. 08/725,070 filed October 2, 1996 (currently pending and now-allowed), which is a continuation-in-part of Application Serial No. 08/407,508, filed March 20, 1995 (now abandoned). The contents of Application Serial Nos. 08/725,070  
5 and 08/407,508, are incorporated herein by reference.

### **FIELD OF THE INVENTION**

The present invention pertains generally to ophthalmic surgery which is useful for correcting vision deficiencies. More particularly, the present invention pertains to methods which surgically correct the vision of a patient  
10 by removing portions of the stroma to reshape the cornea. The present invention is particularly, but not exclusively useful as a method for correcting the vision of a patient by lifting a contoured corneal flap created by a laser beam to expose a bed of stromal tissue, photoaltering the exposed bed of stromal tissue in a predetermined manner and subsequently repositioning the  
15 flap.

### **BACKGROUND OF THE INVENTION**

Vision impairments such as myopia (i.e. near-sightedness), hyperopia (i.e. far-sightedness) and astigmatism can be corrected using eyeglasses or contact lenses. Alternatively, the cornea of the eye can be reshaped  
20 surgically to provide the needed optical correction. For example, it is known that if part of the cornea is removed, the pressure exerted on the cornea by the aqueous humor in the anterior chamber of the eye will act to close the void created in the cornea, resulting in a reshaped cornea. By properly selecting the size, shape and location of a corneal void, the desired shape,  
25 and hence optical properties of the cornea can be obtained.

One procedure employed to reshape the cornea is to remove portions of the anterior portion of the cornea. For example, see U.S. Patent No. 4,665,913 which issued to L'Esperance for an invention entitled "Method for

Ophthalmological Surgery," and U.S. Patent No. 4,669,466 which issued to L'Esperance for an invention entitled "Method and Apparatus for Analysis and Correction of Abnormal Refractive Errors of the Eye." Another procedure used to reshape the cornea removes and reshapes subsurface tissue such as stromal tissue. As an example of such a procedure, U.S. Patent No. 4,907,586, which issued to Bille et al. for an invention entitled "Method for Reshaping the Eye," discloses an intrastromal photoalteration technique for reshaping the cornea. Importantly for the purposes of the present invention, the above cited Bille patent discloses the use of a pulsed laser beam for photoalteration of intrastromal tissue. As disclosed by Bille, the pulsed laser beam penetrates corneal tissue and is focused at a point below the surface of the cornea to photoalter stromal tissue at the focal point. The ability to reach a subsurface location without necessarily providing a physical pathway allows for great flexibility in corneal reshaping and can reduce the total amount of tissue disruption required for a particular corneal reshaping. Further, as the prescribed corneal void shape becomes more complex and precise, the need to access subsurface tissue without a physical pathway becomes more important.

Recently developed so-called LASIK procedures incise the anterior portion of the cornea using a microkerotome to create a flap. It should be recognized that a microkeratome is a mechanical device that uses an automated blade to create a flap. Once created, the flap can be temporarily lifted for photoalteration of the exposed stroma. This procedure, like the procedure disclosed in Bille et al. '586, has as its objective the removal of only stromal tissue with the consequent preservation of anterior corneal tissue. As discussed above, the LASIK procedure relies on a physically prepared pathway, and hence is limited to simple flap configurations. In contrast with the simple flap configurations which can be prepared using a microkerotome, the procedure of the present invention recognizes that a pulsed laser beam can be focused below the surface to create complex flap designs.

In light of the above, it is an object of the present invention to provide a method for corneal laser surgery that corrects the refractive characteristics of

the cornea by removing only stromal tissue and minimizes the total amount of tissue undergoing photoalteration. Another object of the present invention is to provide a method for corneal laser surgery which creates a corneal flap having a complex peripheral edge such as a peripheral edge which can be repositioned in an interlocking relationship with undisturbed corneal tissue to hold the corneal flap in place during subsequent healing, or a peripheral edge that incorporates a tab to assist in lifting and repositioning the corneal flap. Still another object of the present invention is to provide a method for corneal laser surgery which creates a corneal flap that can be lifted to expose and then photoalter a bed of stromal tissue that has a complex shape, such as a convex, concave or irregularly shaped bed. Yet another object of the present invention is to provide a method for corneal laser surgery which is relatively easy to practice and comparatively cost effective.

#### **SUMMARY OF THE PREFERRED EMBODIMENTS**

In accordance with the present invention, a method for corneal laser surgery includes the step of first prescribing the size, shape and location of stromal tissue which needs to be removed in order to correct the vision deficiency of a patient. This volume of stromal tissue which is to be removed is generally in the form of a lentoid that is defined by an anterior surface and a posterior surface and may contain an annular surface. In general, the surfaces of the lentoid can be either plane, convex, concave or irregular. In the method of the present invention, a contoured corneal flap having an interior surface and a peripheral edge is created wherein the interior surface of the flap is shaped to conform to the anterior surface of the prescribed lentoid.

To create the contoured corneal flap, a pulsed laser beam is focused to a preselected start point within the stromal tissue. In accordance with preplanned procedures, the focal point will be located on the intended interior surface of the flap. The focal point is then moved within the stromal tissue to cut (photoalter) a layer of tissue having the desired contour of the interior

surface of the flap (and hence the anterior surface of the prescribed lentoid). Next, the focal point is moved within the cornea to create a peripheral edge for the flap. In this case, the peripheral edge of the flap is a surface that extends from the perimeter of the interior surface of the flap to the anterior surface of the cornea. In the preferred embodiment of the present invention, the peripheral edge may incorporate features which allow the flap to interlock with the cornea when the flap is repositioned. Further, the peripheral edge of the flap may be formed with a tab to assist in lifting and repositioning the flap. Still further, the border of the anterior surface of the flap and the perimeter of the interior surface of the flap, both of which lie on the peripheral edge, are generally curvilinear, but are not closed curves. Rather, the flap is formed with a hinge of corneal tissue which allows for flap rotation about the hinge during lifting and repositioning of the flap relative to the cornea.

Once created, the contoured corneal flap can be lifted to expose a bed of intrastromal tissue. Next, an excimer laser can be used to photoalter the bed of intrastromal tissue in a predetermined manner, thus creating the posterior surface of the prescribed lentoid. Finally, the flap having a contoured inner surface that defines the anterior surface of the lentoid, can be repositioned over the newly created void and allowed to heal. The result is a reshaped cornea that effectively corrects a patient's vision deficiency. As envisioned for the present invention, lasers may be used for plasma mediated tissue ablation (generally superficial tissue) and for plasma mediated tissue disruption (generally internal bulk tissue). Accordingly, the term photoalteration will be used herein to indicate an operation wherein there may be either plasma mediated tissue ablation or plasma mediated tissue disruption.

### **BRIEF DESCRIPTION OF THE DRAWINGS**

The novel features of this invention, as well as the invention itself, both as to its structure and its operation, will be best understood from the accompanying drawings, taken in conjunction with the accompanying



description, in which similar reference characters refer to similar parts, and in which:

Fig. 1 is a perspective view of a patient being treated with a pulsed/excimer laser in accordance with the method of the present invention;

5 Fig. 2 is a perspective view of an eye;

Fig. 3 is a cross sectional view of a portion of the cornea of the eye as seen along the line 3-3 in Fig. 2 showing the anatomical layers of the cornea and a representative lentoid;

Fig. 4 is a plan view of the cornea after the incision of a flap;

10 Fig. 5A is a cross-sectional view of a cornea as seen along the line 5-5 in Fig. 4, showing a flap incision for a flap having a concave interior surface;

Fig. 5B is a cross-sectional view of a cornea as in Fig. 5A, showing the cornea after the incision and lifting of a flap having a concave interior surface;

15 Fig. 5C is a cross-sectional view of a cornea as in Fig. 5B showing the cornea prior to photoalteration of the exposed bed, and showing the posterior and annular surfaces of the lentoid in phantom.

Fig. 5D is a cross-sectional view of a cornea as in Fig. 5C showing the cornea after photoalteration of the exposed bed of stromal tissue;

20 Fig. 5E is a cross-sectional view of a cornea as in Fig. 5D showing the cornea after the removal of a lentoid of stromal tissue from the exposed bed of stromal tissue by photoalteration, and replacement of the flap;

Fig. 5F is a cross-sectional view of a cornea as in Fig. 5E showing the reshaped cornea after removal of a lentoid of stromal tissue;

25 Fig. 6A is a cross-sectional view of a cornea as in Fig. 5A showing the cornea after the incision of a flap having a convex interior surface;

Fig. 6B is a cross-sectional view of a cornea as in Fig. 6A showing the cornea after the incision and lifting of a flap having a convex interior surface;

Fig. 7 is a plan view of a cornea after the incision of an oval flap;

Fig. 8 is a plan view of a cornea after the incision of an elongated flap;

30 Fig. 9 is a plan view of a cornea after the incision of a flap having a tab;

Fig. 10A is a cross-sectional view of a cornea as seen along the line 10-10 in Fig. 9, showing a flap having an integral tab to assist in lifting and repositioning the flap;

Fig. 10B is a cross-sectional view of a cornea as in Fig. 10a, showing the cornea after the incision and lifting of a flap having a tab;

Fig. 11 is a plan view of a cornea after the incision of a flap having an interlocking feature; and

Fig. 12 is a plan view of a cornea after the incision of a flap having a beveled peripheral edge with an acute angle between the peripheral edge and the interior surface of the flap.

### **DESCRIPTION OF THE PREFERRED EMBODIMENT**

Referring initially to Fig. 1, an apparatus 10 for alternately generating either a pulsed laser beam 12 or an excimer laser beam 13 is shown. As hereinafter disclosed in the specification and in Fig. 1, the combined numerals 12/13 will refer respectively to either the pulsed laser beam 12 or the excimer laser beam 13. As contemplated for the present invention, the apparatus 10 will use both laser beams 12/13. Specifically, a pulsed laser beam 12 will first be used to create a flap of corneal tissue and the excimer laser beam 13 will then be used to remove corneal tissue below the flap. It will be appreciated by the skilled artisan that in lieu of an excimer laser, a pulsed infrared laser beam or a visible pulsed laser beam may be used to remove corneal tissue below the flap.

In detail, Fig. 1 shows the pulsed laser beam 12 being directed onto the eye 14 of a patient 16. For purposes of the present invention, a pulsed laser beam 12 preferably has the physical characteristics similar to those of the pulsed laser beams generated by a laser system as disclosed and claimed in U.S. Patent No. 4,764,930, which issued to Josef F. Bille et al. for an invention entitled "Multiwavelength Laser Source." Furthermore, the present invention contemplates the use of a pulsed laser beam 12 which has pulses with durations as long as a few nanoseconds or as short as only a few

femtoseconds. The pulsed laser beam 12 has a fluence of less than 100 joules per square centimeter. Also, as indicated above, the apparatus 10 will generate, a second type of laser beam; namely, an excimer laser beam 13.

Fig. 2 shows the anatomical structure of the human eye 14 including the cornea 18, the pupil 20, the iris 22, and the sclera 24. In Fig. 3 it can be seen that the cornea 18 includes five anatomically definable layers of tissue. Going in a direction from anterior to posterior in Fig. 3, the tissue layers of the cornea 18 are: the epithelium 26, Bowman's membrane 28, the stroma 30, Decemet's membrane 32 and the endothelium 34. Of these, the stroma 30 is of most importance for the present invention as it contains the only tissue which is to be removed for correction of the patient's vision. Also shown in Fig. 3, the anterior chamber 35 is a cavity filled with aqueous humor 37. The pressure exerted by in the aqueous humor 37 maintains the shape of the cornea 18.

As indicated above, it is known that the correction of myopic, hyperopic and astigmatic conditions can be accomplished by the removal of a predetermined volume of stromal tissue 30. Further, the particular volume of stromal tissue 30 to be removed for the prescribed optical correction will depend on the amount and type of correction required and will generally be a lens shaped volume (a lentoid) 36. An example of a lentoid volume 36 is shown in cross-section in Fig. 3. As shown, it is to also be appreciated that the lentoid volume 36 will be defined by an anterior surface 38, a posterior surface 40 and may have a annular surface 39.

In accordance with the methods of the present invention, access to the prescribed lentoid volume 36 is accomplished by using a pulsed laser beam 12 to create a contoured corneal flap 42. By cross-referencing Figs. 4, 5A and 5B, it can be seen that the contoured flap 42 has an interior surface 44 and a peripheral edge 46. A pulsed laser beam 12 is used to create the contoured flap 42 by focusing the pulsed laser beam 12 at a point within the stromal tissue 30 and moving the focal point of the pulsed laser beam 12 within the stromal tissue 30 to cut a subsurface layer 48. Layer 48 is an interface between the interior surface 44 of flap 42 and the bed 50 of stromal

tissue 30, and as such, layer 48 has a shape conforming to the prescribed shape of the interior surface 44 of the flap 42.

Next, the peripheral edge 46 for the flap 42 is created. To create the peripheral edge 46, the pulsed laser beam 12 is focused at a point within the stromal tissue 30 and on the boundary 52 of the bed 50. Then, the focal point of the pulsed laser beam 12 is moved within the stromal 30 to cut a layer 54. Layer 54 extends from the boundary 52 of bed 50 to the anterior surface 56 of the cornea 18. Layer 54 is an interface between the peripheral edge 46 of the flap 42 and the wall 58 that surrounds the bed 50. The points where the peripheral edge 46 of the flap 42 intersects the anterior surface 56 of the cornea 18 is the anterior border 60, and is shown in both Fig. 4 and Fig. 5B. Both the anterior border 60 and the boundary 52 of bed 50 may be curvilinear, but are not necessarily closed curves. Rather, in the preferred embodiment of the present invention, both the boundary 52, and the anterior border 60 terminate within the stroma 30 to create a hinge 62 of stromal tissue 30 for flap 42. Hinge 62 allows the flap 42 to be lifted while continuing to be attached to the remaining cornea 18.

Once the flap 42 is created, the flap 42 can be lifted by rotating the flap 42 about the hinge 62 to expose the bed 50 of stromal tissue 30. The contour of the exposed bed 50 as well as the contour of the interior surface 44 of the flap 42 will conform to the layer 48 cut into the stromal tissue 30 by the pulsed laser beam 12. As shown in Figs. 5C and 5D, after the flap 42 has been lifted and the bed 50 of stromal tissue 30 is exposed, a pulsed laser beam 12 or an excimer laser beam 13 can be used to photoalter a portion or all of the bed 50 in a predetermined manner until the posterior bed surface 64 of stromal tissue 30 is reached. The shape of the posterior bed 64 can be selectively contoured using the laser beam 12,13 to conform to the prescribed shape of the posterior surface 40 of the prescribed lentoid 36, as shown in Fig. 5E. As indicated earlier, lasers may be used for plasma mediated tissue ablation (generally superficial tissue) and for plasma mediated tissue disruption (generally internal bulk tissue). Accordingly, the term photoalteration will be

used herein to indicate an operation wherein there may be either plasma mediated tissue ablation or plasma mediated tissue disruption.

As further shown by cross-referencing Figs. 5D and 5E, after the photoalteration of the prescribed lentoid 36 volume by either an excimer laser beam 13 or a pulsed laser beam 12 is complete, the contoured flap 42 can be reengaged with the cornea 18 into a position covering the lentoid 36. In particular, the flap 42 can be rotated about the hinge 62 until the peripheral edge 46 of the flap 42 is positioned into contact with a portion of the wall 58. When the flap 42 is properly repositioned over the lentoid 36, the anterior surface 56 of the cornea 18 will be smooth and continuous across the anterior border 60 from the flap 42 to the remaining portion of the cornea 18. After repositioning, the flap 42 will heal in place, and this healing will result in a continuous tissue between the peripheral edge 46 of the flap 42 and a portion of the wall 58 of the cornea 18.

Fig. 5E shows the cornea 18 after the flap 42 has been repositioned, and shows an example of a lentoid 36 having an anterior surface 38, an annular surface 39 and a posterior surface 40. Further, Fig. 5F shows the reshaped cornea 18 which results after the methods of the present invention. As discussed above, after a prescribed lentoid 36 of stromal tissue 30 has been removed and the flap 42 repositioned over the lentoid 36, the pressure exerted by the aqueous humor 37 in the anterior chamber 35 will cause the cornea 18 to close the lentoid 36 volume and hence reshape the cornea 18. In particular, the pressure exerted by the aqueous humor 37 will push the posterior bed 64 into contact with the interior surface 44 of the repositioned flap 42, where the two surfaces will subsequently heal together and become continuous stromal tissue 30. By comparing Fig. 5A with Fig. 5F, it can be seen that the curvature of the anterior surface 56 of the reshaped cornea 18 (Fig. 5F) differs from the curvature of the anterior surface 56 of the initial cornea 18 (Fig. 5A).

As can be expected, the lentoid 36 shape shown in Figs. 5A – 5F is only one of the many possible lentoid 36 shapes that can be prescribed and thereafter created by the methods of the present invention. In particular, the

example lentoid 36 shape as shown in Figs. 5A – 5F has a convex anterior surface 38, a concave posterior surface 40 and an annular surface 39 connecting the anterior 38 and posterior 40 surfaces. As shown, the contour of the convex anterior surface 38 does not necessarily have the same curvature as the anterior surface 56 of the cornea 18. Rather, the points on the layer 48 cut by the pulsed laser beam 12 are located at variable distances from corresponding points on the anterior surface 56 of the cornea 18. Although not required by the method of the present invention, the lentoid 36 may have anterior 38 and posterior 40 lentoid surfaces that have the same approximate curvature, such as the lentoid 36 shown in Figs. 5A – 5F. When this type of lentoid 36 is prescribed, it can be conveniently created using an excimer laser 13 configured to photoalter the exposed bed 50 of stromal tissue 30 to a uniform depth.

Figs. 6A and 6B show an example of a flap 42 that can be cut using the methods of the present invention to create a prescribed lentoid 36 having a concave anterior surface 38. As discussed above, the versatility of the pulsed laser beam 12, alone or in combination with an excimer laser beam 13, enables one skilled in the art to create a flap 42 in accordance with the present invention which will result in a lentoid 36 having a plane, concave, convex or irregularly shaped anterior surface 38, and a plane, concave, convex or irregularly shaped posterior surface 40.

Further, as shown in Fig. 7, using the methods of the present invention, an oval flap 68 can be created having an oval anterior border 70. One benefit of the oval shape for flap 68 is that the oval shape allows for a bed 50 with a large exposed bed area. Similarly, as shown in Fig. 8, an elongated flap 72, having an elongated anterior border 74 can be created with the methods of the present invention. An elongated flap 72 may also provide the benefit of exposing a bed 50 with a large exposed bed area.

Additionally, custom shaped flaps 76 can be created using the methods of the present invention. For example, as shown by cross-referencing Figs. 9 and 10A-B, a custom flap 76 having a tab 66 can be made. Referring to Fig. 9, the tab 66 may have a different curvature than the custom anterior border

78 of the flap 76, and hence the tab 66 extends from the custom anterior border 78 to assist in lifting and repositioning the custom flap 76.

In accordance with the methods of the present invention, an interlocking flap 80 as shown in Fig. 11 can be created for the purposes of maintaining the flap 80 in place after repositioning to both facilitate healing and reduce any optical distortions that may occur if a repositioned flap 42 shifts before healing is completed. As shown in Fig. 11, the interlocking flap 80 contains an interlocking peripheral edge 82. In one embodiment of the interlocking peripheral edge 82, an annular ring 84 extends from the interlocking peripheral edge 82 for engagement with a corresponding recess 86 formed in the wall 88.

Fig. 12 shows an alternative embodiment of an interlocking flap 90, having a beveled peripheral edge 92 for interlocking of the flap 90 with the remaining cornea 18 after repositioning. In the embodiment shown in Fig. 12, the flap 90 is formed with the angle  $\alpha$  between the beveled peripheral edge 92 and the interior surface 44 of the flap 90 as an acute angle. A flap 90 with a beveled peripheral edge 92 as shown in Fig. 12 is further disclosed in co-pending and now-allowed Application No. 08/725,070 entitled "Method for Corneal Laser Surgery," which is incorporated herein by reference.

While the particular Method of Corneal Reshaping by Laser Incising a Contoured Corneal Flap as herein shown and disclosed in detail is fully capable of obtaining the objects and providing the advantages herein before stated, it is to be understood that it is merely illustrative of the presently preferred embodiments of the invention and that no limitations are intended to the details of the construction or design herein shown other than as defined in the appended claims.

What is claimed is:

1. A method for reshaping the cornea of an eye under a flap of corneal tissue which comprises the steps of:

5 cutting a subsurface layer of stromal tissue, said subsurface layer being an interface between a interior surface for said flap and a bed of stromal tissue, said interior surface located on said flap opposite a portion of the anterior surface of the cornea and at a selected distance therefrom;

10 incising the cornea to create a peripheral edge for said flap of corneal tissue, said interior surface of said flap being bounded by said peripheral edge;

lifting said flap to expose said bed;

photoaltering at least a portion of said bed of stromal tissue to correct the visual acuity of the eye; and

15 repositioning said flap over said bed.

2. A method as recited in claim 1 wherein said selected distance is variable to create a convex shape for said interior surface of said flap.

3. A method as recited in claim 1 wherein said selected distance is variable to create a concave shape for said interior surface of said flap.

20 4. A method as recited in claim 1 wherein said bed of stromal tissue has a boundary and said boundary is substantially in the shape of a circle.

25 5. A method as recited in claim 1 wherein said bed of stromal tissue has a boundary and said boundary is substantially in the shape of a oval.



6. A method as recited in claim 1 wherein said incising step and said cutting step are accomplished using a pulsed laser beam.

7. A method as recited in claim 6 wherein said photoaltering step is accomplished using an excimer laser.

5 8. A method as recited in claim 1 wherein said peripheral edge of said flap borders said portion of the anterior surface of the cornea and said peripheral edge of said flap has a substantially uniform predetermined depth.

9. A method as recited in claim 1 wherein said peripheral edge of said flap is formed with a tab to assist in lifting and repositioning of said flap.

10 10. A method as recited in claim 1 wherein said peripheral edge of said flap is formed with an interlocking feature to hold said flap in place after said repositioning step.

11. A method as recited in claim 1 wherein said photoaltering step is accomplished using a pulsed infrared laser.

12. A method for reshaping the cornea of an eye under a flap of corneal tissue which comprises the steps of:

focusing the rays of a pulsed laser beam within the stroma of the cornea to photoalter stromal tissue at only said focal point;

5 moving said focal point of said pulsed laser beam along a predetermined path within the stroma of the cornea to photoalter a layer of stromal tissue having a preselected shape, said layer being an interface between the interior surface of said flap and a bed of stromal tissue;

10 incising the cornea between the anterior surface of the cornea and the preselected layer to create a peripheral edge for said flap, said flap substantially overlying said bed of stromal tissue;

lifting said flap to expose said bed of stromal tissue;

15 photoaltering at least a portion of said bed of stromal tissue to create a void in the stromal tissue of the cornea; and replacing said flap over said void.

13. A method as recited in claim 12 wherein said void is lens-shaped having an anterior surface, a posterior surface and an annular surface.

20 14. A method as recited in claim 13 wherein said anterior surface is concave shape.

15. A method as recited in claim 13 wherein said posterior surface is convex shape.

25 16. A method as recited in claim 13 wherein said anterior surface is convex shape.

17. A method as recited in claim 12 wherein said bed of stromal tissue has a boundary and said boundary is substantially in the shape of a circle.

18. A method as recited in claim 12 wherein said bed of stromal tissue has a boundary and said boundary is substantially in the shape of a oval.

19. A method as recited in claim 12 wherein said photoaltering step is accomplished using an excimer laser.

20. A method as recited in claim 12 wherein said peripheral edge of said flap is formed with a tab to assist in lifting and repositioning of said flap.

21. A method as recited in claim 12 wherein said peripheral edge of said flap is formed with an interlocking feature to hold said flap in place after said repositioning step.

22. A method as recited in claim 12 wherein said photoaltering step is accomplished using a pulsed infrared laser.

23. A method as recited in claim 12 wherein said photoaltering step is accomplished using a visible pulsed laser.

## **ABSTRACT**

In accordance with the present invention, a method for corneal laser surgery includes the step of first prescribing the size, shape and location of a lentoid of stromal tissue to be removed in order to correct the vision of a patient. Next, a contoured corneal flap having an interior surface and peripheral edge is created wherein the interior surface of the flap is shaped to conform to the prescribed anterior surface of the lentoid. To create the flap, the focal point of a pulsed laser beam is moved within the intrastromal tissue to photoalter a layer of tissue in the shape of the interior surface and the peripheral edge of the flap. Once created, the flap can be lifted to expose a bed of intrastromal tissue. Then, an excimer laser can be used to photoalter the bed of intrastromal tissue in a predetermined manner, thus creating a void in the shape of the prescribed lentoid. Finally, the flap can be repositioned over the lentoid shaped void and allowed to heal. The result is a reshaped cornea that effectively corrects a patient's vision impairment.

PATENT: 11236.11

03/27/00 4:14 PM




Fig. 1

Fig. 2




Fig. 3

